

ANTI-INFLAMMATORY ACTIVITY OF *IN VITRO* AND *IN VIVO* *KIGELIA PINNATA* (JACQ.) DC FROM INDIAN ORIGIN

Purnima Kumari¹, Reena Sachan², Poonam Yadav¹, Parul Tomer¹, Arvind Arya³, Sandeep Tripathi¹ and *Sandeep Kumar¹

¹Department of Biotechnology, NIET, Nims University Jaipur, Jaipur, Rajasthan, India

²Department of Biotechnology, Uttaranchal college of Science and Technology, Dehradun, India

³Department of Biotechnology, Meerut institute of Engineering and Technology, Meerut, U.P., India

*Author for Correspondence

ABSTRACT

Anti-inflammatory activities of methanolic extract of callus and intact plant part of *Kigelia pinnata* (Jacq.) DC (Bignoniaceae) was studied. Acute inflammation model, carrageenan induced rat hind paw edema model was employed to investigate the anti-inflammatory activity. The methanolic extract from fresh callus cultures of *Kigelia pinnata* cultured on Murashige and Skoog (MS) medium fortified with α -NAA (1.0 mg/l) and BAP (0.05 mg/l) for 2 - 12 weeks was studied and compared with intact plant extract. Results from the anti-inflammatory activity testing indicated that the callus extract showed a less significant inhibition in the edema as compared to extract of intact plant. It was also revealed that there is an increase in the accumulation of secondary metabolites in callus cultures of sixth week of incubation. The methanolic extract of callus culture and intact plant part exhibited significant, dose-dependent activity on the tested experimental animal models and produced a significant inhibition of carrageenan-induced paw edema in rat. The study suggests the plant *Kigelia pinnata* is a potent source for phytomedicine development in future.

Key Words: Callus Culture, Carrageenan, Paw Edema, Inflammation, *Kigelia pinnata*. (Jacq.) DC, In Vitro, In Vivo, Medicinal Plant

INTRODUCTION

Inflammation is a very complex, multifactorial and dynamic process involving many systems which is closely associated with the process of repair. Inflammation is defined as local response of living mammalian tissue to injury due to any agent and manifests usually in form of painful swelling associated with some changes in skin covering the site. Inflammation can be classified as either as acute or chronic. Acute inflammation is the initial response of the body to harmful stimuli and is due to the increased movement of plasma and leukocytes from the blood into the injured tissues. Chronic inflammation is due to a progressive shift in the type of cells which are present at the site of inflammation which is characterized by simultaneous destruction and healing of the tissue from the inflammatory process. Carrageenan-induced paw edema model is widely used for determining the acute phase of the inflammation. Histamine, 5-hydroxytryptamine and bradykinin are the first detectable mediators in the early phase of carrageenan-induced inflammation whereas prostaglandins are detectable in the late phase of inflammation (Panchal *et al.*, 2011; Jain *et al.*, 2011).

Plant based traditional medicine system continues to play an essential role in health care with about 80% of world's inhabitants relying on it for their primary health care (Garcia *et al.*, 2003). *Kigelia pinnata* (Jacq.) DC is a woody plant that belongs to Bignoniaceae family. It is also known as Balam kheera, Jhar fanoos and Sausage tree because of its sausage shaped fruits. It is distributed throughout tropical regions of the world. In India it is abundantly found in West Bengal, U.P, Rajasthan etc. It is habitant of waterlogged areas like wet savannah and riverine areas (Gragg and Newman 2001; Sofowora 1982). The adult tree attains a height up to 20 m length. The bark is with greyish colour, smooth and peeling texture

Research Article

at early and late stages respectively. Flowers are in panicle with reddish to orange or purplish green colour.

The sausage tree is being used by ancient times for its medicinal properties. The economical important parts of the tree include fruit, bark, root and leaves. Along with its medicinal uses it is also a functional component of beverage, cosmetics industries and animal husbandary (livestock green fodder). The plant extract is reported to have molluscicidal, antibacterial, antifungal (Binutu *et al.*, 1996), antiprotozoal (Hoet *et al.*, 2004; Moideen *et al.*, 1999), antineoplastic (Inoue *et al.*, 1981) and anticarcinogenic effects. It is also useful in diarrhoea (Akah 1998), toothache (Gill, 1992), ringworm (Binutu 1996), Dysentery (Bharti *et al.*, 2006) etc. An anti-malarial compound known as lapachol (Hussain *et al.*, 2007) has been extracted from the root of *K. pinnata* as marvelous alternate to chloroquine and quinine.

Conventionally, *Kigelia pinnata* (Jacq.) DC reproduces via viable seeds, but the low percentage of seed viability limits its natural propagation. Hence alternative methods like micropropagation through tissue culture could open the door for its cultivation with pure line breeds in short span of time.

The investigation of certain indigenous plants for their anti-inflammatory activities can play an important role in drug development program in the pharmaceutical industries.

The aim of this study was to investigate plant and its callus to determine their anti-inflammatory activity and to determine whether the ethno botanical approach in finding anti-inflammatory is a useful approach.

MATERIALS AND METHODS

Plant Materials

Young leaves and shoots of *Kigelia pinnata* (Jacq.) DC were used as explant to initiate callus culture and the leaves were used for the extract preparation.

Drugs and Chemicals

The drugs and chemicals used were Carrageenan (Sd Fine Chemicals Limited, Mumbai), Gum acacia, Indomethacin (Indocap, Jagsonpal Pharmaceuticals Ltd., Faridabad) and Methanol (BDH, Mumbai). Murashige and Skoog medium (Himedia Laboratories Pvt. Ltd., Mumbai), Agar Type 1 (Central Drug House Pvt. Ltd. New Delhi), Sucrose (RFCL Ltd., New Delhi), α -Naphthalene acetic acid (Central Drug House Pvt. Ltd., New Delhi) 2,4,5-Trichlorophenoxy acetic acid (Himedia Laboratories Pvt. Ltd., Mumbai) and 6-Benzylaminopurine (Sigma, China).

Animals

A permission from Institutional Animal Ethics Committee was granted and after wards male Wistar rats (150-250 g) were maintained under standard husbandry conditions and acclimatized to the laboratory environment for a period of one week prior to the experimental session. All the animals were divided into different groups each consists of six animals. The experimental animal models were made to fast overnight prior to the experiments.

Callus Culture Conditions

Young leaves and stem of *Kigelia pinnata* (Jacq.) DC rinsed with distilled water and were sterilized in 70% ethanol for 1 min and then followed by five times washing with sterile distilled water and explants were aseptically cut in segments of 0.5 cm length. These explants were cultured in sterile polystyrene flasks containing 40 ml of MS medium (Luyindula *et al.*, 2004). The medium was fortified with a mixture of α -NAA, 1.0 mg/l and BAP, 0.05 mg/l prior to autoclaving at 121°C (1.5 Kg/cm²) for 30 min, the pH of all media was adjusted to 5.8 with 0.1N NaOH. All cultures were incubated at 25°C with photoperiod for 16 h and darkness for 8 h in a culture room. The initiated callus were routinely sub-cultured onto a fresh (multiplication) medium fortified with same concentration of growth regulators as in initiation medium to proliferate the callus at faster rate.

Measurements

Fresh weights of the calli were recorded immediately after the cultivation time and dried weights were determined by drying to a constant weight at 60°C in an oven for 24 h.

Research Article

Preparation of Extracts

The plant leaves were shade dried, powdered and extracted with methanol while callus were dried in oven at 35°C and extracted with methanol. The extracts were filtered and concentrated and kept in a vacuum desiccator for complete removal of the solvent. Methanol extract of leaves and calli were obtained in the yield of 11.07% and 60.7% w/w respectively. The extracts were stored at 4°C in glass vials for use in anti-inflammatory activities.

Phytochemical Screening

The methanolic extract was examined for the presence of various phyto-constituents like alkaloids, tannins, flavonoids, phenolics and glycosides by employing standard phytochemical tests (Parekh and Chanda, 2006).

Anti-inflammatory Activity

The animals were divided into six groups. Group I (n=6) served as Control, received the vehicle only (1% CMC, 10 ml/kg oral dose). Group II (n=6) served as Standard, received Indomethacin at dose of 1 mg/kg body weight. Group III (18 animals in subgroups of 6), Group IV (18 animals in subgroups of 6) served as test, received methanolic extract of *Kigelia pinnata* (Jacq.) DC leaves (KPL) and *Kigelia pinnata* (Jacq.) DC callus (KPC) at oral doses of 100, 200 and 400 mg/kg body weight respectively.

Paw Edema Induction

Carrageenan induced paw edema was used to determine the anti-inflammatory activity of the extracts (Perez *et al.*, 1990). The animals pretreated with extract or Indomethacin one hour before were injected with 0.1 ml of 1% Carrageenan (in 1% CMC) solution into the sub-plantar region of right hind paw. Paw volume was measured by displacement of the water column in a Plethysmometer (Ugo Basile, Italy) immediately after Carrageenan application at 0, 1, 2, 3, 4 and 6 hrs after the stimulus. Reduction in the paw volume compared to the vehicle-treated control animals was considered as anti-inflammatory response.

The percentage inhibition of edema was calculated as follows:

$$\text{Percentage of Inhibition} = \frac{(V_T - V_O) \text{ Control} - (V_T - V_O) \text{ Treated group}}{(V_T - V_O) \text{ Control}} \times 100$$

V_O = paw volume of the rat before administration of Carrageenan .

V_T = paw volume of the rat after the administration of Carrageenan at different time intervals.

Percentage inhibition of paw edema was found to be proportional to anti-inflammatory activity.


RESULTS AND DISCUSSION

MS medium fortified with α -NAA (1 mg/l) and BAP (0.05 mg/l) was found best for initiation and multiplication of callus from leaf and shoot explant. The callus cultures were maintained in dark conditions in a culture room. For further multiplication the callus were transferred on to the multiplication medium. The physical appearance of all calli was friable and ranging from yellow to brown (Kumar *et al.*, 2003, 2004). Figure 2-4 shows that the growth of different callus cultures (2, 4, 6, 8, 10 and 12 weeks) expressed as the increase or decrease in fresh and dry weights. The weights of fresh callus extracts significantly increased with the duration of cultivation and then decreases. The dried callus weight was maximum of sixth week of callus culture. The maximum secondary metabolites were also seen in the sixth week incubation of callus culture. After extraction, the extracts were subjected to qualitative phytochemical tests (Table 1).


The anti-inflammatory effect of the extract and the reference drug in carrageenan induced paw edema model in rats is shown in Table 2. Carrageenan induced paw edema in rats reached to a peak value at 4 hrs and graded doses of methanolic extract of plant and callus produced a significant inhibition in the edema volume ($P < 0.001$). The leaf extract at the test doses 100, 200 and 400 mg/kg body weight reduced the edema induced by carrageenan by 28, 37 and 42% respectively at 4 h and callus extract

Research Article


reduced the edema by 25, 28 and 34% respectively at 4h, whereas the standard drug showed 85% of inhibition as compared to the control group. All the doses of methanolic extract of KP(P) are significantly inhibit the paw edema in rats than all the doses of KP(C). The study suggests the methanolic extracts of intact plant part of *Kigelia pinnata* has more potent inhibition than callus culture for carrageenan induced paw edema in rat.




Initiation of callus from leaf and stem explant (2 weeks)



Multiplication of callus culture (4 and 6 weeks old culture)



Multiplication of callus culture (8 and 10 weeks old culture)



Multiplication of callus culture (12 weeks old culture)

Figure 1: Callus cultures of *Kigelia pinnata* (Jacq.) DC on MS medium supplement with α -NAA (1.0 mg/l) + BAP (0.05 mg/l)




Figure 2: Effect of culturing period on fresh weight of callus on MS medium with α -NAA and BAP




Figure 3: Effect of culturing period on dry weight of callus




Figure 4: Effect of culturing period on percentage loss of biomass of callus on drying

Table 1: Qualitative phytochemical test of extracts of KP(P) and KP(C)

S. No	Test/Reagent used	Sample KP(P)	Sample KP(C)
Test for alkaloids			
1	Mayer's reagent	+ ve	+ve
2	Dragendorff's reagent	+ ve	+ ve
Test for Carbohydrates and Glycosides			
5	Molish's reagent	+ ve	+ ve
6	Fehling's solution	+ ve	- ve
Test for Tannins and phenolic compounds			
7	Ferric chloride solution	+ ve	+ ve
8	Geletin solution	+ ve	- ve
9	Lead acetate solution	+ ve	+ ve
Test for lignin			
10	Extract + Alcoholic solutions of phloroglucinol and hydrochloric acid	+ ve	- ve
Test for Terpenes			
11	Extract + chloroform+ acetic anhydride + Sulfuric acid	+ ve	+ ve
Test for Flavonoids			
12	Methanolicextract + Hydrochloric acid + Magnesium ribbon	+ ve	+ ve

Carrageenan induced paw edema is widely used for determining the acute phase of the inflammation (Jain *et al.*, 2011) and is characterized by biphasic event with involvement of different inflammatory mediators. In the first phase (during the first 2 hrs after carrageenan injection), chemical mediators such as histamine, dextran (Das *et al.*, 2010) and serotonin play role, while in second phase (3-4 hrs after carrageenan injection) Kinin and prostaglandins are involved (Hernandez and Rabanal 2002; Ratheesh and Helen, 2007). The results of present study revealed that administration of methanolic extract inhibited the edema starting from the first hour and during all phases of inflammation, which probably leads to inhibition of different aspects and chemical mediators of inflammation.

It is well known that in chronic and sub-acute inflammation reactive ROS play an important role in modulating the extent of inflammatory response and consequent tissue and cell injury (Garcia *et al.*, 2003) and antioxidants are considered as possible protective agents reducing oxidative damage of human body from ROS and retard the progress of many diseases (Ngemenya *et al.*, 2006). The natural phenolic, alkaloids, tannins, glycosides and flavonoids compounds function as antioxidants by different mechanisms and according to present study, the high contents of these phytochemicals in both the extracts can explain its anti-inflammatory activity (Bairagi *et al.*, 2012).

Table 2: Anti-inflammatory effect of the extract and reference drug in carrageenan induced paw edema in rats

Groups	Dose mg/kg b.w	Paw Edema Volume							
		1hr		2hrs		3hrs		4hrs	
		Mean \pm S.E.M	%PEI	Mean \pm S.E.M	%PEI	Mean \pm S.E.M	%PEI	Mean \pm S.E.M	%PEI
Control	----	0.30 \pm 0.01	-----	0.85 \pm 0.02	-----	0.90 \pm 0.01	----	0.50 \pm 0.01	----
Standard	10	0.25 \pm 0.03	33	0.30 \pm 0.02 ^c	78	0.30 \pm 0.01 ^c	80	0.20 \pm 0.01 ^c	85
KP(P)	100	0.23 \pm 0.04	13	0.64 \pm 0.05 ^b	22	0.65 \pm 0.04 ^c	26	0.35 \pm 0.03 ^b	28
	200	0.17 \pm 0.02 ^c	20	0.55 \pm 0.04 ^c	28	0.56 \pm 0.04 ^c	32	0.27 \pm 0.03 ^c	37
	400	0.26 \pm 0.03	26	0.64 \pm 0.03 ^c	30	0.63 \pm 0.02 ^c	36	0.35 \pm 0.03 ^b	42
KP(C)	100	0.18 \pm 0.02 ^b	13	0.63 \pm 0.03 ^c	17	0.66 \pm 0.02 ^c	18	0.31 \pm 0.04 ^b	25
	200	0.27 \pm 0.02	20	0.68 \pm 0.06 ^a	24	0.70 \pm 0.06 ^a	26	0.40 \pm 0.04	28
	400	0.21 \pm 0.03 ^a	26	0.60 \pm 0.04 ^c	28	0.61 \pm 0.03 ^c	32	0.33 \pm 0.04 ^b	34

Values are expressed as mean \pm S.E.M; n=6; significance at $p < 0.05^a$, 0.01^b and 0.001^c as compared to the control. PEI – Paw Edema Inhibition

Thus, it can be concluded that the methanolic extract of *in vivo* plant possess higher anti-inflammatory activity than leaf and shoot callus. Further studies involving the purification of the chemical constituents of the plant and the investigations in the biochemical pathways may result in the development of a potent anti-inflammatory agent with low toxicity and better therapeutic index.

REFERENCES

- Akah PA (1998).** Anti diarrheal activity of *Kigelia africana* in experimental animals. *Journal of Herbs, Spices and Medicinal Plants* 31-38.
- Bairagi SM, Aher AA, Nema N and Nimase PK (2012).** Anti-inflammatory evaluation of methanol extraction and aqueous fraction of the bark of *Bauhinia variegata* (Leguminosae). *International Journal of Research in Pharmacy and Chemistry* 2(1) 77-82.
- Bharti N, Singh S, Fehmida N and Amir A (2006).** Isolation and *in vitro* antiamoebic activity of iridoids isolated from *Kigelia pinnata*. *Archive for Organic Chemistry* 69-76.
- Binutu OA, Adesogan KE and Okogun JI (1996).** Antibacterial and antifungal compound from *Kigelia pinnata*. *Planta Medica* 62(4) 352-353.
- Das S, Haldar PK, Goutam P and Suresh RB (2010).** Evaluation of anti-inflammatory activity of *Clerodendron infortunatum* Linn. extract in rats. *Global Journal of Pharmacology* 4(1) 48-50.
- Garcia D, Leiro J, Delgado R, Sanmartin ML and Ubeira FM (2003).** *Mangifera indica* L. extract (Vimang) and mangiferin modulate mouse humoral immune responses. *Phytotherapy Research* 17 1182–1187.
- Gill LS (1992).** Ethnomedical uses of plants in Nigeria. Uniben Press, Benin city p. 143.
- Gragg GM and Newman DJ (2001).** Medicinal for the millennia: the historical record. *Annals of the New York Academy of Sciences* 95(3) 3-25.

Research Article

Hernandez PM and Rabanal Gallego R (2002). Evaluation of the anti-inflammatory and analgesic activity of *Sideritis anariensis* var. *pannosa* in mice. *Journal of Ethnopharmacology* **81** 43-47.

Hoet S, Oppendoes F, Brun R and Quetin-Leclercq J (2004). Natural products active against African trypanosomes: a step towards new drugs. *Natural Product Report* **21** 353-364.

Hussain H, Krohn K, Ahamad UV, Miana GA and Green IR (2007). Lapachol: an overview. *Archive for Organic Chemistry* **2** 145-171.

Inoue K, Inoue H and Chen C (1981). Quinones and related compounds in higher plants, A naphthoquinone and a lignin from the wood of *Kigelia pinnata*. *Phytochemistry* **20** 2271-2276.

Jain S, Garg V K and Sharma P K (2011). Anti-Inflammatory Activity of Aqueous Extract of *Beta Vulgaris* L. *Journal of Basic and Clinical Pharmacy* **2**(2) 83-86.

Kumar S, Singh KK and Rai LK (2004). In vitro propagation of a endangered Sikkim Himalayan rhododendron (*R.maddeni*) from cotyledonary nodal segments. *Journal of American Rhododendron Society* **58**(2) 101-105.

Kumar S, Suri SS, Sonie KC and Ramawat KG (2003). Establishment of embryogenic cultures and somatic embryogenesis in callus cultures of guggul (*Commiphora wightii* Arnott. Bhandari). *Indian Journal of Experimental Biology* **41** 69-77.

Luyindula N, Tona L, Lunkebila S, Tsakala M, Mesia K, Musuamba CT, Cimanga RK, Apers S, De Bruyne T, Pieters L and Vlietinck AJ (2004). In Vitro Antiplasmodial Activity of Callus Culture Extracts from Fresh Apical Stems of *Phyllanthus niruri*: Part 1. *Pharmaceutical Biology* **42**(7) 512-518.

Moideen SVK, Houghton PJ, Rock P, Croft SL and Aboagye-Nyame F (1999). Activity of extract and naphthoquinones from *Kigelia pinnata* (Jacq.) DC against *Trypanosoma brucei brucei* and *Trypanosoma brucei rhodesiense*. *Planta Medica* **65** 493-588.

Murashige T, Skoog F (1962). A revised media for rapid growth and bioassays with Tobacco tissue cultures. *Physiological Plantarum* **15** 473-497.

Ngemenya MN, Mbah JA, Tane P and Titanji VPK (2006). Antibacterial effects of some Cameroonian medicinal plants against common pathogenic bacteria. *African Journal of Traditional, Complementary and Alternative Medicines* **3** 84-93.

Panchal A H, Patel R K, Pundarikakshudu K and Bhandari A (2011). Evaluation of anti inflammatory activities of extract of leaves of *Eclipta alba* and *Widelia calandulaceae*. *An International Journal of Pharmaceutical Science* 1846-1852.

Parekh J and Chanda S (2006). In vitro antimicrobial activities of extracts of *Launaea procumbens* Roxb. (Labiateae), *Vitis vinifera* L. (Vitaceae) and *Cyperus rotundus* L. (Cyperaceae). *African Journal Biomedical and Research* **9** 89-93.

Perez C, Pauli M and Bazevque P (1990). An antibiotic assay by the agar well diffusion method. *Acta Biologiae et Medicinae Experimentalis* **15** 113-115.

Ratheesh M and Helen A (2007). Anti-inflammatory activity of *Ruta graveolens* Linn. On carrageenan induced paw edema in wistar male rats. *African Journal of Biotechnology*. **6**(10) 1209-1211.

Sofowora A (1982). Medicinal plants and traditional medicine in Africa. Published by John wiley and Sons Ltd 1st Edn. **131** 168-171.