

Research Article

**ANTIGENIC CHARACTERIZATION OF EXCRETORY / SECRETORY
ANTIGEN OF *OESOPHAGOSTOMUM COLUMBIANUM* USING
WESTERN BLOTTING**

***S. Arunkumar**

*Department of Veterinary Parasitology, Madras Veterinary College
Tamilnadu veterinary and animal sciences university, Chennai-600007, India*

**Author for Correspondence*

ABSTRACT

In the present investigation, E/S antigen from *O. columbianum* was isolated by *in vitro* culture method and hyper immune serum was raised against this antigen in rabbits. On SDS – PAGE analysis, characterization, the E/S antigen revealed six polypeptides at 29.0, 38.0, 49.0, 71.0, 84.0 and 95.0 kDa. Further, three immuno-reactive polypeptides at 29.0, 38.0 and 49.0 kDa were detected in western blotting. This antigenic characterization study will be useful in developing suitable sero diagnostic in Oesophagostomosis in sheep and goats.

Key Words: *Oesophagostomum Columbianum, E/S Antigen, Western Blotting, Goats*

INTRODUCTION

Oesophagostomum columbianum is a common and widely prevalent nematode with significant economic importance in small ruminants (Olivares *et al.*, 2001). Caprine oesophagostomosis is insidious and chronic in nature and the migratory prepatent stages of the parasite are involved in its major pathogenic effect. Analysis of host immune responses to helminthes is hampered by two main factors (i) the complexity of antigenic profiles (ii) the presence of cross- reactive determinants on antigens. To identify specific antigens, excretory/secretory antigens have received increasing attention recently. This is due to the fact that E/S antigens relatively display less complexity compared to somatic antigens. Further, the work on antigenic composition of E/S antigen of *O. columbianum* is scanty in India. Hence, the present study was undertaken to analyze the polypeptide profiles of E/S antigen of *O. columbianum* for its use in serodiagnostic or immuno protective aspect.

MATERIALS AND METHODS

Preparation of E/S antigen

Adult live *Oesophagostomum columbianum* worms were collected from the colon of goats slaughtered at corporation slaughter house, Perambur, Chennai. The collected worms were washed five times in normal saline and subsequently washed five times in Phosphate buffered saline (PBS, pH 7.4), containing penicillin (500 IU/ml) and streptomycin (5mg/ml) and nystatin (1mg/100ml). Then, the worms were identified based on morphological features using standard keys (Soulsby, 1982).

The fresh and highly motile worms were transferred to RPMI 1640 medium containing penicillin (100 IU/ 100 ml) and streptomycin (1mg/100ml) and cultured at a concentration of approximately 400 worms per 20 ml in a culture flask at 5 per cent CO₂ atmosphere at 37°C for 24 hours. The medium was changed every 6 hours after incubation and fresh medium was added with 2 per cent glucose throughout incubation. Worm viability was monitored throughout this period on the basis of motility, integrity of the worms. Moreover, random samples of the culture fluid obtained during and directly after the incubation period were plated out on agar in order to exclude bacterial contaminations. After the incubation period, the culture medium was collected by decantation and filtered through a 0.22 µm filter (Millipore). Then, the culture medium was centrifuged at 10,000 rpm for 30 minutes at 4°C and the supernatant was labelled as excretory/ secretory (E/S) antigen. This culture procedure was repeated several times in order to obtain

Research Article

sufficient quantity of antigen. Finally, the antigen obtained was concentrated by dialysis (membrane cut off, 12 kDa) against polyethylene glycol (PEG 6000 – SRL, India) over a period of 6 hours. The concentrated material was added with 1 mM phenyl methyl sulfonyl fluoride (PMSF) and 1 mM ethylene diamine tetra acetic acid (EDTA) to prevent proteolysis and stored at -20°C with 0.02 per cent sodium azide as preservative till further use. The protein concentration of the E/S antigen was determined using bicinchoninic acid (BCA) method (Smith *et al.*, 1985) using protein estimation kit (Genei, Bangalore).

Raising of Hyper Immune Serum

New Zealand white rabbits were immunized by intramuscular injection with E/S antigen (0.5mg) emulsified with Freund's complete adjuvant. Subsequently, three booster doses of antigen, emulsified with Freund's incomplete adjuvant were given at weekly intervals. Blood from the hyperimmunized rabbits was collected after 7 days of the last booster. Serum was separated and stored at -20° C for using in the assay.

Characterization of Antigen

Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) was carried out to observe the polypeptide patterns of E/S antigen of *O. columbianum* according to their molecular weights under reducing gel conditions as per method of Laemmli (1970). The mini gel (12%) electrophoresis (Biorad, USA) was performed using 1mm thickness gel in a discontinuous system. The denatured antigen sample was loaded into each well with protein marker in a separate well. The gel was electrophoresed at a constant voltage of 120 till the tracking dye reached one cm above the lower extremity of the gel. After electrophoretic run, the gel was subjected to Coomassie brilliant blue (HiMedia, India) staining overnight with gentle agitation. After proper staining, the gel was put in destaining solution until clear demarcation of bands appeared.


The immuno-reactive polypeptides were identified by western blotting technique (Towbin *et al.*, 1979) using the hyper immune serum raised in rabbits. After the electrophoretic run, the nitrocellulose membrane was removed and was blocked in blocking buffer at 37°C for 2 hours. Then the membrane was washed in washing buffer three times with gentle agitation for 5 minutes each. The membrane was incubated in immune serum at a dilution of 1:100 in PBS for 2 hours at 37°C with gentle shaking. The nitrocellulose membrane was washed in washing buffer three times for 5 minutes each and immersed in solution of 1:1000 diluted anti-sheep IgG HRP conjugate (Sigma, USA) in PBS for 2 hours at 37°C. Then the membrane was washed three times in washing buffer for 5 minutes each. The membrane was treated with substrate solution (DAB) till the bands appeared.

RESULTS AND DISCUSSION


During incubation period, adult worms remained viable as assessed qualitatively by both motility and clumping tendency. No bacterial contamination was detected during *in-vitro* culture. The protein concentration of the E/S antigen was found to be 1.8 mg/ml. Similar culture procedures were reported for isolating the E/S antigen by other workers (Anbu and Joshi, 2008; Karanu *et al.*, 1993). The protein concentration obtained in this study is in accordance with the reports of Jas *et al.*, (2010).

On Characterization the E/S antigen revealed six polypeptide bands at 29.0, 38.0, 49.0, 71.0, 84.0 and 95.0 kDa molecular weights by SDS-PAGE analysis (Figure-1). Out of six polypeptides, three polypeptides were identified as low molecular weights (29.0, 38.0, 49.0 kDa) and another three were identified as medium range molecular weights (71.0, 84.0, and 95.0 kDa). On western blot analysis, the E/S antigen probed with rabbit hyper immune serum showed three immuno-reactive polypeptides at 29.0, 38.0 and 49.0 kDa molecular weights (Figure-2). A very limited literature is available on antigenic profiles of *O. columbianum* for comparison. Jas *et al.*, (2010) analysed the antigenic composition of E/S antigen of *O. columbianum* by SDS- PAGE and observed ten polypeptides with molecular weights ranged between 22.5 to 98.0 kDa. Further, they demonstrated two immuno-reactive polypeptides at 39.0 and 50.0 kDa molecular weights in western blotting.

Research Article



M – Molecular weight marker
L1, L2 – E/S antigen of *O. columbianum*
Figure -1: SDS-PAGE analysis of E/S antigen of *Oesophagostomum columbianum*



M – Molecular weight marker
L1 – E/S antigen of *O. columbianum*
Figure -2: Western blot analysis of E/S antigen of *O. columbianum* against homologous hyper immune serum

In recent years, E/S products of helminths have received increasing attention. This is due to the fact that E/S antigens usually display a relatively simple antigenic composition compared to the somatic antigen. Further, they also play numerous role in the host parasite interactions. The sensitivity and specificity of E/S antigens were high compared to somatic antigens in immuno assays (Mir *et al.*, 2008). Hence, the antigenic composition of E/S antigen of *O. columbianum* was analysed in this study.

Summary

In the present study, E/S antigen from *O. columbianum* was prepared and hyper immune serum was raised against this antigen in rabbits. On antigenic characterization, the E/S antigen revealed six polypeptides at 29.0, 38.0, 49.0, 71.0, 84.0 and 95.0 kDa. On western blotting, three immuno-reactive polypeptides at 29.0, 38.0 and 49.0 kDa were detected. This study will be helpful in developing suitable serodiagnostic in Oesophagostomosis in sheep and goats.

Research Article

REFERENCES

- Anbu KA and Joshi P (2008).** Identification of a 55 kDa *Haemonchus contortus* excretory / secretory glycoprotein as a neutrophil inhibitory factor. *Parasite Immunology* **30** 23-30.
- Jas R, Ghosh JD and Das K (2010).** Poly clonal antibody based coproantigen detection immuno assay for diagnosis of *Oesophagostomum. columbianum* infection in goats. *Veterinary Parasitology* **170** 262-267.
- Karanu FN, Rurangirwa FR, McGuire TC and Jasmer DP (1993).** *Haemonchus contortus*: Identification of proteases with diverse characteristics in adult worm excretory – secretory products. *Experimental Parasitology* **77** 362-371.
- Laemmli UK (1970).** Cleavage of structural proteins during assembly of the head of Bacteriophage T4. *Nature* **227** 680-685.
- Mir RA, Chishti MZ, Zargar MA and Ganie SA (2008).** Excretory / Secretory antigen are better than crude antigen for serodiagnosis of *Haemonchus contortus*. *Asian Journal of Scientific Research* **1**(2) 171- 175.
- Olivares JL, Rodriguez DJ, Herrera H, Cortes S and Gonzalez O (2001).** Experimental *Oesophagostomum columbianum* infection in ovines. *Revista de Salud Animal* **23** 118- 121.
- Smith PK, Krohn RI, Hermanson GT Mallia AV and Klenk D C (1985).** Measurement of protein using bicinchoninic acid. *Analytical Biochemistry* **150** 76-85.
- Soulsby E.J.L. (1982).** Helminths, Arthropods and Protozoa of domesticated animals. 7th edition Bailliere Tindall, London 186.
- Towbin H, Stachelin T and Gordon J (1979).** Electrophoretic transfer of proteins from polyacrylamide gels to nitro cellulose sheep: procedure and some application. *Proceedings of National Academy of Sciences USA* **76** 4530- 4534.