

## **EVALUATION OF ABSOLUTE GROWTH RATE IN SHORT TERM SELECTION FOR DIFFERENT AGES IN JAPANESE QUAIL**

**\* A. Ashok<sup>1</sup> and R. Prabakaran<sup>2</sup>**

*Department of Poultry Science, Madras Veterinary College, TANUVAS, Chennai,  
Tamil Nadu, India*

*\*Author for Correspondence*

*Present address: <sup>1</sup>Institute of Food and Dairy Technology, Tamilnadu Veterinary and Animal Sciences  
University, Koduvalli, Alammathi Post, Chennai, Tamilnadu, India.*

*<sup>2</sup>Tamilnadu Veterinary and Animal Sciences University, Chennai, Tamilnadu, India*

*\*Author for Correspondence*

### **ABSTRACT**

Meat type Japanese quail were subjected to three different methods of individual phenotypic selection viz., high two week body weight, four week body weight and high four week body weight coupled with low relative body weight gain between 4-6 weeks of age. Selection was carried out for three generations and the respective populations were designated as Line selected for high two week body weight (SWL), Line selected for high four week body weight (FWL), Line selected for high four week body weight and low relative weight gain between 4-6 weeks of age (LWL) lines and a control line (COL) was also maintained without adopting any selection with random pairing. The overall least squares means for absolute growth rate (AGW) during 1,2,3,4,5 and 6 weeks of age were  $26.31 \pm 0.31$ ,  $37.79 \pm 0.54$ ,  $39.76 \pm 0.58$ ,  $31.93 \pm 0.59$ ,  $21.28 \pm 0.52$  and  $18.37 \pm 0.48$  g, respectively. The least squares means of AGW ranged from  $26.96 \pm 0.35$  g at first week to  $37.42 \pm 0.69$  g third week and dropping steeply to  $19.01 \pm 0.53$  g at sixth week in SWL line. In FWL, AGW varied from  $28.03 \pm 0.34$  at 1<sup>st</sup> week,  $40.93 \pm 0.59$  g at 2<sup>nd</sup> week to  $19.59 \pm 0.52$  g at 6<sup>th</sup> week. AGW in LWL was  $25.10 \pm 0.36$  at 1<sup>st</sup> week  $49.08 \pm 0.66$  g at 3<sup>rd</sup> week and  $17.43 \pm 0.55$  g at 6<sup>th</sup> week and the corresponding values for COL were  $25.13 \pm 0.36$ ,  $45.57 \pm 0.61$  and  $17.44 \pm 0.55$  g, respectively. The findings indicated that selection for high body weight over generations might have shifted the peak in the growth curve to early ages.

**Key Words:** *Japanese quail- selection- body weights- absolute growth rate*

### **INTRODUCTION**

In India, commercial quail farming has created a huge impact in recent years and many farms have been established throughout the country both for meat and egg production. Japanese quails are marketed at 4-5 weeks of age for meat with a body weight of 150-180 g. The females come into lay at about 6-7 weeks of age, reared for breeding upto 6-8 months of age and produce about 90-150 eggs with a weight of about 8-13 g. Tamilnadu is growing into a major producer and consumer of Japanese quail meat in India.

The growth rate of quail chicks is normally quite rapid from hatch to five weeks of age and slows down thereafter. Unlike all other domestic avian species, sexual dimorphism favours the female body weight in quail. Experimental research established that body weight of Japanese quail responded quickly to selection (Nestor *et al.*, 1982; Caron *et al.*, 1990 and Marks, 1993). Selection for rapid growth at a specific age has resulted in a tremendous growth response. With this marked response there has been concomitant decrease in age at which body weight selection is applied. Thus, within a relatively short period of time, selection age and market weights have shifted to earlier ages and heavier weights respectively (Anthony, 1995).

A selection experiment was designed. Individual phenotypic selection was contemplated to facilitate development of superior breeder flock suitable for production of optimum number of fast growing

## Research Article

commercial meat type Japanese quails. The study was also designed to obtain an understanding of the relationship between selection age and growth with the following objectives, viz., to evaluate selection for juvenile, fourth-week and sixth-week body weights in Japanese quail.

### MATERIALS AND METHODS

The study was carried out at the Institute of Poultry Production and Management, formerly known as Poultry Research Station, Tamilnadu Veterinary and Animal Sciences University, Nandanam, Chennai, India. A Japanese quail (*Coturnix japonica*) population, maintained at the institute formed the base population for this study. The foundation stock for the three selected and an unselected control populations was from a random mating Japanese quail line maintained at the Institute of Poultry Production and Management, Chennai. The line had no known history of artificial selection except for a short period during 1989 to 1992 when the population was subjected to selection on the basis of body weight at four weeks of age for four generations under two different nutritional environments of high and low protein diets. From the foundation stock, one hundred and eighty males and equal number of females were randomly selected, wing banded, weighed, and randomly assigned to four groups to have 45 pairs in each of the four groups. The breeder males and females were maintained in cages under single pair mating. Hatching eggs were collected and set for hatch. Chicks hatched from three groups were subjected to individual phenotypic selection for body weight at different ages. One group (SWL) was selected for high body weight at two weeks of age, the other (FWL) for high body weight at four weeks of age. The third group (LWL) was subjected to two stage selection with the initial selection practised at four weeks of age for high body weight, followed by selection for low relative body weight gain between four to six weeks of age. The fourth group (COL) was maintained as control line with random selection of parents.

The number of hatches obtained and the total number of progenies produced in the three selected lines and control were 2176, 1780, 2331 and 2343, respectively in  $S_0$ ,  $S_1$ ,  $S_2$  and  $S_3$  generations. Only those data of progenies with intact wing bands and whose sexes were phenotypically identifiable were included in the study. One of the four groups formed in the base generation ( $S_0$ ) was treated as control line and raised separately along with the selected populations (other three lines) in each generation to observe and account for environmental influences. Single pair mating was followed with females assigned at random to individual males with the restriction that no full sib mating was permitted.

### Measurement of the traits

**Body weight gain:** The data of progenies whose wing bands were intact, sexes were identifiable and also only those with body weight records from hatch to six weeks of age were included in the study.

**Absolute body weight gain (AGW):** The absolute body weight gain (AGW) was computed by the following formula as suggested by Kratochvílová *et al.* (2002).

$$\text{Absolute body weight gain} = \frac{(\text{Body weight 2} - \text{Body weight 1})}{(\text{Age 2} - \text{Age 1})}$$

The AGW during 1<sup>st</sup> (AGW1), 2<sup>nd</sup> (AGW2), 3<sup>rd</sup> (AGW3), 4<sup>th</sup> (AGW4), 5<sup>th</sup> (AGW5) and 6<sup>th</sup> (AGW6) week of age were computed based on the difference in body weights between two successive intervals.

## Research Article

**Statistical analysis:** The data generated on body weight for age were corrected for the fixed effects of line, generation, sex and hatch by the least squares analysis (Harvey, 1979) using the following linear model based on pooled data.

$$Y_{ijklm} = \mu + st_i + g_j + s_k + h_l + e_{ijklm}$$

Where,

$Y_{ijklm}$  = measurement of a trait on  $m^{th}$  bird belonging to  $l^{th}$  hatch,  $k^{th}$  sex,  $j^{th}$  generation and  $i^{th}$  line

$\mu$  = overall mean

$st_i$  = effect of  $i^{th}$  line

$g_j$  = effect of  $j^{th}$  generation

$s_k$  = effect of the  $k^{th}$  sex

$h_l$  = effect of  $l^{th}$  hatch

$e_{ijklm}$  = random error, assumed to be distributed normally and independently with mean zero and variance  $\sigma^2$

Duncan's multiple range test (Duncan, 1955) was employed to make all pair wise comparisons of means.

## RESULTS AND DISCUSSION

The results of least squares analysis of variance of AGW based on pooled data presented in Table- 1. The overall least squares means for AGW during 1, 2,3,4,5 and 6 weeks of age were  $26.31 \pm 0.31$ ,  $37.79 \pm 0.54$ ,  $39.76 \pm 0.58$ ,  $31.93 \pm 0.59$ ,  $21.28 \pm 0.52$  and  $18.37 \pm 0.48$  g, respectively (Table - 2).

The least squares means of AGW ranged from  $26.96 \pm 0.35$ g at first week to  $37.42 \pm 0.69$  g third week and dropping steeply to  $19.01 \pm 0.53$  g at sixth week in SWL line. In FWL, AGW varied from  $28.03 \pm 0.34$  at 1<sup>st</sup> week,  $40.93 \pm 0.59$  g at 2<sup>nd</sup> week to  $19.59 \pm 0.52$  g at 6<sup>th</sup> week. AGW in LWL was  $25.10 \pm 0.36$  at 1<sup>st</sup> week  $49.08 \pm 0.66$  g at 3<sup>rd</sup> week and  $17.43 \pm 0.55$  g at 6<sup>th</sup> week and the corresponding values for COL were  $25.13 \pm 0.36$ ,  $45.57 \pm 0.61$  and  $17.44 \pm 0.55$  g, respectively.

**Table: 1 Least squares analysis for variance of absolute weight gain**

	Lines		Generations			Sexes		Hatches		Error
	d.f	M.S.S	d.f	M.S.S	d.f	M.S.S	d.f	M.S.S	d.f	M.S.S
<b>AGW1</b>	3	2773.38**	3	6795.13**	1	519.87**	5	25745.89**	5549	49.72
<b>AGW2</b>	3	5899.62**	3	26820.34**	1	2297.48**	5	9290.66**	5549	149.17
<b>AGW3</b>	3	3294.73**	3	115484.44**	1	9362.21**	5	45551.47**	5549	173.19
<b>AGW4</b>	3	6472.73**	3	27437.96**	1	27455.05**	5	49921.60**	5549	177.66
<b>AGW5</b>	3	3329.69**	3	8542.22**	1	80789.21**	5	22244.90**	5549	134.53
<b>AGW6</b>	3	1625.29**	3	353.17*	1	153470.61**	5	2924.08**	5549	114.74

\*Significant at  $P < 0.05$ ; \*\* Significant at  $P < 0.01$

**Table: 2 Least squares means of absolute body weight gain (g) at various ages (0-6 weeks) on pooled data**

	AGW1			AGW2			AGW3			AGW4			AGW5			AGW6		
	N	Mean	SE	N	Mean	SE	N	Mean	SE	N	Mean	SE	N	Mean	SE	N	Mean	SE
<b>Overall</b>	<b>5562</b>	<b>26.31</b>	<b>0.31</b>	<b>5562</b>	<b>37.79</b>	<b>0.54</b>	<b>5562</b>	<b>39.76</b>	<b>0.58</b>	<b>5562</b>	<b>31.93</b>	<b>0.59</b>	<b>5562</b>	<b>21.28</b>	<b>0.52</b>	<b>5562</b>	<b>18.37</b>	<b>0.48</b>
<b>Line</b>																		
SWL	1469	26.96 <sup>b</sup>	0.35	1469	37.19 <sup>b</sup>	0.60	1427	37.42 <sup>c</sup>	0.69	1427	34.36 <sup>a</sup>	0.66	1427	22.36 <sup>a</sup>	0.57	1427	19.01 <sup>a</sup>	0.53
FWL	1307	28.03 <sup>a</sup>	0.34	1307	40.93 <sup>a</sup>	0.59	1263	26.96 <sup>d</sup>	0.70	1263	33.12 <sup>a</sup>	0.65	1263	22.86 <sup>a</sup>	0.56	1263	19.59 <sup>a</sup>	0.52
LWL	1443	25.10 <sup>c</sup>	0.36	1443	36.42 <sup>b</sup>	0.62	1541	49.08 <sup>a</sup>	0.66	1541	29.96 <sup>b</sup>	0.68	1541	20.30 <sup>b</sup>	0.59	1541	17.43 <sup>b</sup>	0.55
COL	1343	25.13 <sup>c</sup>	0.36	1343	36.62 <sup>b</sup>	0.63	1331	45.57 <sup>b</sup>	0.61	1331	30.30 <sup>b</sup>	0.69	1331	19.60 <sup>b</sup>	0.60	1331	17.44 <sup>b</sup>	0.55
<b>Generation</b>																		
S <sub>0</sub>	1427	26.38 <sup>c</sup>	0.37	1427	43.66 <sup>a</sup>	0.64	1427	37.42 <sup>b</sup>	0.69	1427	27.70 <sup>b</sup>	0.70	1427	23.76 <sup>a</sup>	0.61	1427	18.02 <sup>c</sup>	0.56
S <sub>1</sub>	1263	22.99 <sup>d</sup>	0.37	1263	32.93 <sup>d</sup>	0.65	1263	26.96 <sup>c</sup>	0.70	1263	27.82 <sup>b</sup>	0.71	1263	21.27 <sup>b</sup>	0.61	1263	17.83 <sup>c</sup>	0.57
S <sub>2</sub>	1541	28.44 <sup>a</sup>	0.36	1541	36.59 <sup>c</sup>	0.62	1541	49.08 <sup>a</sup>	0.66	1541	36.24 <sup>a</sup>	0.67	1541	17.91 <sup>c</sup>	0.58	1541	18.57 <sup>bc</sup>	0.54
S <sub>3</sub>	1331	27.42 <sup>b</sup>	0.33	1331	37.97 <sup>b</sup>	0.57	1343	37.60 <sup>b</sup>	0.68	1343	35.99 <sup>a</sup>	0.62	1343	22.17 <sup>b</sup>	0.54	1343	19.06 <sup>a</sup>	0.50
<b>Sex</b>																		
Male	2774	26.00 <sup>a</sup>	0.33	2774	37.15 <sup>b</sup>	0.57	2774	38.46 <sup>b</sup>	0.62	2774	29.71 <sup>b</sup>	0.62	2774	17.46 <sup>b</sup>	0.54	2774	13.11 <sup>b</sup>	0.50
Female	2788	26.61 <sup>a</sup>	0.32	2788	38.43 <sup>a</sup>	0.56	2788	41.06 <sup>a</sup>	0.60	2788	34.16 <sup>a</sup>	0.61	2788	25.10 <sup>a</sup>	0.53	2788	23.63 <sup>a</sup>	0.49
<b>Hatch</b>																		
1	1634	25.86 <sup>c</sup>	0.18	1634	35.88 <sup>d</sup>	0.30	1634	37.94 <sup>b</sup>	0.33	1634	45.79 <sup>a</sup>	0.33	1634	26.41 <sup>b</sup>	0.29	1634	20.04 <sup>b</sup>	0.27
2	1638	20.50 <sup>e</sup>	0.18	1638	42.27 <sup>b</sup>	0.30	1638	51.64 <sup>a</sup>	0.33	1638	29.59 <sup>d</sup>	0.33	1638	28.07 <sup>a</sup>	0.29	1638	18.75 <sup>c</sup>	0.27
3	1515	22.33 <sup>d</sup>	0.18	1515	38.54 <sup>c</sup>	0.32	1515	38.58 <sup>b</sup>	0.34	1515	33.92 <sup>c</sup>	0.35	1515	20.41 <sup>c</sup>	0.30	1515	16.17 <sup>e</sup>	0.28
4	708	36.73 <sup>a</sup>	0.28	708	34.91 <sup>c</sup>	0.49	708	36.30 <sup>c</sup>	0.53	708	38.10 <sup>b</sup>	0.54	708	14.55 <sup>e</sup>	0.47	708	17.34 <sup>d</sup>	0.43
5	42	17.49 <sup>f</sup>	1.11	42	29.22 <sup>f</sup>	1.92	42	51.31 <sup>a</sup>	2.07	42	15.01 <sup>c</sup>	2.09	42	18.35 <sup>d</sup>	1.82	42	13.44 <sup>f</sup>	1.68
6	25	34.9 <sup>b</sup>	1.43	25	45.92 <sup>a</sup>	2.48	25	22.79 <sup>a</sup>	2.67	25	29.20 <sup>d</sup>	2.71	25	19.88 <sup>d</sup>	2.36	25	24.48 <sup>a</sup>	2.17

### Research Article

Means with different superscripts within each column, trait and effect differ significantly ( $P < 0.05$ ). Least squares means of absolute body weight gain over all ages indicated that the same grew rapidly up to two weeks, steadied at 3<sup>rd</sup> week and declined every week thereafter up to six weeks. Contrarily, Sharma and Panda (1978) registered a peak during fourth week while Panda *et al.* (1980) registered the same during third week of age of Japanese quail. Aggrey and Cheng (1994) reported almost similar weight gains during second, third and fourth week of age with the later being marginally higher.

The findings indicated that selection for high body weight over generations might have shifted the peak in the growth curve to early ages.

### REFERENCES

**Aggrey SE and Cheng KM (1994).** Animal model analysis of genetic (co)variance of growth traits in Japanese quail. *Poultry Science* **73** (12) 1822-1828.

**Anthony NB (1995).** Selection for increased body weight in poultry. What role does selection age have on response? Center of Excellence for Poultry Science, Department of Poultry Science, University of Arkansas, Fayetteville Arkansas. Available: <http://www.poultryscience.org/pba/1952-2003/1995/1995%20Anthony.PDF> [Accessed 10 January 2010]

**Caron N, Minivelle F, Desmarad M and Poste LM (1990).** Mass selection for 45-day body weight in Japanese quail: selection response, carcass composition, cooking properties and sensory characteristics. *Poultry Science* **69** (7) 1037-1045.

**Duncan D B (1955).** Multiple range and multiple F tests. *Biometrics* **11** (1) 1-42.

**Harvey WR (1979).** Least squares analysis of data with unequal sub-class numbers. *USDA, Agricultural Research Service*.

**Kratochvílová M, Hyanková L, Knížetová H, Fiedler J and Urban F (2002).** Growth curve analysis in cattle from early maturity and mature body size viewpoints. *Czech J. Anim. Science* **47** (4) 125-132.

**Marks H L (1993).** Carcass composition, Feed Intake and Feed efficiency following L ong -Term selection for Four week body weight in Japanese quail. *Poultry Science* **72** (6) 1005-1011.

**Nestor KE, Bacon WL and Lambio AL (1982).** Divergent selection for body weight and yolk precursor in *Coturnix coturnix Japonica*. 1. Selection response. *Poultry Science* **61** (1) 12-17.

**Panda B, Ahuja SD, Prakashbabu M and Gulati DP (1980).** Evaluation of a quail line for some important economic traits. *Indian Journal of Animal Sciences* **50** (6) 518-520.

**Sharma GL and Panda B (1978).** Studies on some productive traits in Japanese quail (*Coturnix coturnix japonica*). *Indian Poultry Gazette* **62** (1) 24-30.