

ANTIMICROBIAL SCREENING OF SOME TREE SPECIES OF JALORE DISTRICT OF RAJASTHAN

***B.B.S. Kapoor and Manju Saharan**

Herbal Research Laboratory, P.G. Department of Botany, Dungar College, Bikaner 334001, India

**Author for Correspondence*

ABSTRACT

Antimicrobial screening of ethyl ether and alcoholic leaf extracts of three selected tree species growing in Jalore district of Rajasthan was carried out *Cordia dichotoma*, *Cordia gharaf* and *Dichrostachys cinerea* showed positive reactions against bacterial pathogens i.e. *Staphylococcus aureus*, *Escherichia coli* and a fungal pathogen *Candida albicans*.

Keywords: Antimicrobial Screening, Tree Species, Jalore District, Rajasthan

INTRODUCTION

The tree species growing in Jalore district of Rajasthan are a potential source of secondary metabolites of pharmaceutical interest like flavonoids, sterols, steroidal sapogenins, alkaloids, phenolic compounds, sulphides, isothiocyanates, anthocynins, terpenoids etc. These are the active principles, which act as antioxidants, anticarcinogenic, antimicrobials and immunity stimulants.

From this region of Rajasthan, three medicinal tree species of *Cordia dichotoma*, *Cordia gharaf* and *Dichrostachys cinerea* have been selected for antimicrobial screening.

A number of plants have been screened for their antimicrobial activity (Khan *et al.*, 1996; Akhtar *et al.*, 1997; Akhtar *et al.*, 1997; Turkusay and Onogal, 1998; Ahmed *et al.*, 1999; Khan *et al.*, 2001; Ahmed and Beg, 2001).

The antimicrobial principles and their distribution in plants have been reported by many workers (Kapoor and Kumar, 2005; Kapoor *et al.*, 2007; Kapoor *et al.*, 2011; Kapoor and Mishra, 2012; Kapoor and Mishra, 2013; Kapoor and Purohit, 2013; Kapoor and Pandita, 2013; Kapoor and Kumar, 2014; Kapoor and Swami, 2015).

MATERIALS AND METHODS

Present investigation describes the antimicrobial activities of leaf extracts of three selected tree species against Bacterial pathogens *Staphylococcus aureus* (Gram positive), *Escherichia coli* (Gram negative) and *Candida albicans* (Fungal pathogen).

Fresh leaves of all the selected tree species were collected from study area and pulverized into a paste. Cold extraction was done by blending the paste with ethyl ether and 50% ethanol in the ratio of 1: 2, in a Warring Blender at 2500 rpm for 10 min.

The mixture was centrifuged at 3000 rpm. The supernatant was evaporated to dryness and the residue was suspended in double distilled water.

The micro-organisms used for screening were *Staphylococcus aureus* (Gram positive), *Escherichia coli* (Gram negative) and *Candida albicans* (Fungal pathogen). The growth medium used for *Staphylococcus aureus* and *Escherichia coli* was Nutrient broth (10% peptone, 0.5% labanco and 0.5% NaCl, pH adjusted to 7.5) and for *Candida albicans* Sabourands liquid medium (1% peptone, 4% glucose, pH adjusted to 5.8).

Paper discs of known concentration of standard antibiotics namely chloramphenicol, penicillin and mycostatin were used for comparison. Blank paper discs were used as control.

Control discs dipped in ethyl ether and 50% ethanol; plates (5 each for *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*) were employed for each extract. The ratio of inhibition zone the various test samples was compared with the inhibition zone from the high concentration antibiotic reference discs (Khanna *et al.*, 1971).

Research Article

RESULTS AND DISCUSSION

Antimicrobial screening of ethyl ether and alcoholic (50% ethanol) leaf extracts of *Cordia dichotoma*, *Cordia gharaf* and *Dichrostachys cinerea* showed positive reactions against all the three test organisms. (Table 1)

Table 1: Antimicrobial Screening of Leaf Extracts of Selected Tree Species and Standard Reference Antibiotics

Plants	Leaves Extracts	Test Organisms				
		<i>E. Coli</i>		<i>S. Aureus</i>		<i>C. Albicans</i>
		I/C ^a	I/P ^a	I/C ^a	I/S ^a	I/m ^a
<i>Cordia Dichotoma</i>	Ether Extract	1.02	1.46	0.65	0.66	0.59
	Alcoholic Extract	0.53	0.69	0.65	0.64	0.60
<i>Cordia Gharaf</i>	Ether Extract	0.62	0.95	1.36	1.12	0.66
	Alcoholic Extract	1.24	1.75	0.82	0.68	0.68
<i>Dichrostachys Cinerea</i>	Ether Extract	1.15	0.84	0.93	0.89	0.74
	Alcoholic Extract	0.94	0.78	0.82	0.75	0.82

a= Ratio of diameters of the inhibition zone to extracts (10 µg) under observation (I) and diameter of inhibition zone due to standard reference antibiotics.

C= Chloramphenicol (30 µg) against *S. aureus* = 30 mm and *E. coli* 32 mm.

P= Penicillin (10 units) against *S. aureus* = 32 mm.

S= Streptomycin (10 µg) against *E. coli* = 20 mm.

M= Mycostatin (100 units) against *C. albicans* = 32 mm.

Maximum antibacterial activity was exhibited by the extracts of leaves (ether extract and alcoholic extract) of *Cordia gharaf* against *Escherichia coli* and *Staphylococcus aureus* whereas leaf extracts of *Dichrostachys cinerea* shows maximum activity against *Candida albicans*.

Conclusion

The present study indicates that these tree species growing in this Jalore region of Rajasthan have definitely some antimicrobial principles as secondary products, which are responsible for antibacterial and antifungal activity. Thus, the activity of all these test extracts against both bacterial and fungal pathogens, indicate that these arid plants are more resistant to bacterial and fungal attacks due to the presence of some biologically active substances. So, these tree species can be used in pharmaceutical and drug industries.

ACKNOWLEDGEMENT

We express sincere thanks to Head, Department of Microbiology, S.P. Medical College, Bikaner for providing research facilities.

REFERENCES

- Ahmed I & Beg-Arina Z (2001).** Antimicrobial and Phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *Journal of Ethnopharmacology* **74**(2) 113-123.
- Ahmed-El-Sawi S Abd-El-Megeed HF & Ali AM (1999).** Flavonoid and antimicrobial volatiles from *Adhatoda vasica* Nees. *Pharmaceutical and Pharmacological Letters* **9**(2) 52-56.
- Akhtar M, Rahber A, Bhati MH & Aslam M (1997).** Antimicrobial activity of plant decoctions against *Xanthomonas campestris* pv. Citri. *International Journal of Pest Management* **43**(2) 149-153.
- Akhtar M, Rahber A, Bhati MH & Aslam M (1997).** Antimicrobial activity of plant decoctions against *Xanthomonas campestris* pv. Citri. on detached citrus leaves. *Tropical Agriculture* **74**(3) 226-228.

Research Article

- Kapoor BBS and Kumar S (2005).** Herbal plants of Rajasthan Desert; A potential source of antimicrobial principles. *Journal of Arid land Studies Japan* **16** 425- 426.
- Kapoor BBS and Kumar S (2014).** Antimicrobial Screening of Some Medicinal Plants of Barmer District of Rajasthan. *Universal Journal of Pharmacy* **3**(5) 17 – 19.
- Kapoor BBS and Mishra R (2013).** Antimicrobial Screening of Some Arid Herbal Plants of Rajasthan. *Universal Journal of Pharmacy* **02**(01) 110 – 112.
- Kapoor BBS and Mishra R (2013).** Antimicrobial Screening of Some Capparidaceous Medicinal Plants of North-west Rajasthan. *Indian Journal of Pharmaceutical and Biological Research* **1**(2) 20 – 22.
- Kapoor BBS and Pandita S (2013).** Antimicrobial Screening of Some Exotic Tree Species of Rajasthan Desert. *Indian Journal of Pharmaceutical and Biological Research* **1**(3) 8-11.
- Kapoor BBS and Purohit V (2013).** Antimicrobial Screening of Some Fabaceous Plant Species of Rajasthan Desert. *International Journal of Pharmacy and Biological Sciences* **3**(3) 201-203.
- Kapoor BBS and Swami S (2015).** Antimicrobial Activities of Some Medicinal Tree Species of Pali District of Rajasthan. *CIBTECH Journal of Microbiology* **4**(3) 44 – 46.
- Kapoor BBS, Acharya S and Mishra R (2012).** Antimicrobial Screening of Some Medicinal Plants of the Rajasthan Desert: *International Proceedings of Chemical, Biological and Environmental Engineering*, ICBECE, 2012 held at Bangkok, Thailand on 24-25 Nov **46** 105-109.
- Kapoor BBS, Acharya S, Mishra R and Swami S (2011).** Antimicrobial Screening of Some Arid Tiliaceous Plants. *Journal of Phytological Research* **24**(2) 255 – 256.
- Kapoor BBS, Bhumika and Khatri JS (2007).** Antimicrobial activities of medicinal tree species of Hanumangarh District of Rajasthan. *Journal of Phytological Research* **20**(2) 325- 326.
- Khan AJ, Zouba AA & Seapy DG (1996).** Antifungal activity from leaves of *Acacia nilotica* against *Pythium aphanidermatum*. *Journal for Scientific Research, Agricultural Sciences* **1** 7-11.
- Khan MR, Kihara M & Omolosa AD (2001).** Antimicrobial activity of *Castanopsis accuminatissima*. *Fitoterapia* **72**(2) 174-176.
- Khanna P, Mohan S and Nag TN (1971).** Antimicrobials from plant tissue cultures. *Lloydia* **34** 168-170.
- Turkusay B & Onogal E (1998).** Studies on antifungal effects of some plant extracts *in vitro*. *Turkish Journal of Agriculture and Forestry* **22**(3) 267-271.