

## ISOLATION AND CHARACTERIZATION OF ACTIVE SECONDARY METABOLITES FROM SOME ETHNOMEDICINAL PLANTS OF RAJASTHAN

**\*Aparna Pareek**

Department of Botany  
University of Rajasthan,  
Jaipur-302004

\*Author for Correspondence: [aparna992000@yahoo.com](mailto:aparna992000@yahoo.com)

### ABSTRACT

Survey was conducted in the south east region of Rajasthan to study the ethno medicinal plants used by various indigenous people in curing their health ailments. Structured questionnaire was used for the same. The ethno medicinal plants selected were used for photochemical analysis and further for antimicrobial studies based on the presence of active principles. The stem and leaf part of these plants showed antimicrobial activity.

**Keywords:** Secondary Metabolites, Ethnomedicine, Rajasthan

### INTRODUCTION

Ethnomedicinal plants are the most important source of life saving drugs for the majority of world's population. It is estimated that around 80% of people world-wide rely chiefly on traditional, largely herbal medicines to meet their primary health care needs and have gained renewed interest for various reasons, affordability, low pricing, no side effects, their solutions for dreadly diseases and disorders time tested remedies and preventive approaches. Over the years such medicines have gained upward trend for consumption especially with the development and standardization of herbal medicines (Malik *et al.*, 2008).

Plant derived drugs have a long history in both indigenous and modern system of medicine as herbal remedies or crude drugs, or as purified compounds approved by the Food and Drug Administration and similar regulatory agencies. According to one estimate 20,000 to 35,000 species of these plants are used as medicines, pharmaceuticals, cosmetics and nutraceuticals by different indigenous groups all over the world. In most of these plant species active constituents are exploited in modern medicines and referred to as plants of scientific knowledge.

Increase in the market demands has posed threats to phytoresources due to unscrupulous mode of collections. The old method of cultivation of some of the ethnomedicinal plants is relatively expensive and production of medicinal compounds can be elicited *in vitro*. Due to overexploitation of medicinal plants for medicine and scientific research, many of them are facing danger of extinction; therefore it is imperative to adopt alternative methods for rapid propagation.

In the recent scenario, tissue culture has emerged as a promising technique to obtain genetically pure elite populations under *in vitro* conditions. Plant secondary metabolites are economically important as drugs, fragrances, pigments, food additives and pesticides. Secondary metabolites could be enhanced with the help of bioreactors and has a tremendous potential for the large scale synthesis of therapeutically active compounds in medicinal plants. The major advantages of plant cell culture system over the conventional cultivation of plants are: (1) Important compounds can be produced under controlled conditions independent of climatic conditions; (2) Cultured cells would be microbes free (3) Automated control of cell growth and rational regulation of metabolite processes would reduce labour costs. (4) Organic substances are easily extractable from callus cultures.

According to Alferman *et al.* (2003), studies have been carried to use plant cell culture for commercial production of plant secondary metabolites. Four commercially viable *in vitro* secondary metabolite production systems have been created including shikonin (*Lithospermum erythrorhizon*), ginsenosides

## Research Article

(*Panax ginseng*), purpurin (*Rubia akane*), and paclitaxel (*Taxus spec.*). Alferman et al. (2003) further studied that the lack of significant application was due to factors including a lack of storage cells for accumulation of these bioactive compounds. In addition, Preil (2005), blamed lack of progress specifically on the high cost of bioreactors, slow growth of the plants, and low yields of active metabolites. Taxol, paclitaxel, is a compound with anticancer properties that was originally extracted from the leaves and bark of plants of the genus *Taxus*. This compound is being successfully produced via cell suspension culture by Phyton Biotech (Germany) in 70,000 L bioreactors (Wink *et al.*, 2005). In most cases, however, callus culture failed to produce significant quantities of medicinal compounds because of the need for compartmentalization of cellular synthetic processes which require differentiation for proper pathway functionality (De Luca and St Pierre, 2000). Immobilization of such suspension cultures was originally conceived as a possible enhancement of the system to allow the productive cells to be maintained while the bioactive compounds were removed with majority of the media (Baldi *et al.*, 2007). Immobilization of *Capsicum* cell cultures allowed for continual production of capsaicin in vitro (Johnson *et al.*, 1990).

The inherent variability in the phytochemical composition of herbal compounds as demonstrated by Bauer and Tittle (1996) can be controlled through the use of culture techniques. *In vitro* culture systems can also be used for specific regulation of medicinal compounds for production of higher value materials.

## MATERIALS AND METHODS

The ethno medicinal plants were collected from the south east area of Rajasthan during the survey conducted. The plants were collected on the basis of their use in curing various health ailments of the indigenous people residing in that particular area. The plants were further identified from the Herbarium in Department of Botany, University of Rajasthan, Jaipur (India). Following ethnomedicinal plants i.e *Argeria speciosa*, *Soymida febrifuga*, *Peganum harmala*, *Celastrus paniculata*, *Gmelina arborea*, *Oogenia sps*, *Gymnema sylvestris* depending on the presence of active constituents in them were used for further antimicrobial studies. Following methods were employed for successful extraction of phytochemicals in these ethnomedicinal plants.

### Extraction and Isolation of Secondary Metabolites

#### (i) Extraction:-

Ethanol was selected as initial extraction solvent due to its ability to dissolve a vast variety of compounds in it. Literature also showed that polar extracts also possessed all biological activity. Plant powder/Callus biomass will be soaked in ethanol in appropriate container and will be regularly shaken for six days at room temperature ( $28 \pm 2^\circ\text{C}$ ) in which solvent will be collected and replaced every day (Ghisalberti, 2000). Then it was filtered through Whatmann filter paper No. 45. Filtrate was dried under vacuum evaporator at  $40^\circ\text{C}$  to yield concentrated dry extract. This extract was stored at  $4^\circ\text{C}$  for further investigations.

#### (ii) Isolation and purification using Column Chromatography and Thin Layer Chromatography (TLC):-

(a) **First column:** Stationary phase used was silica gel with mesh number of 230-400 and the mobile phase was 20% ethyl acetate/hexane. Dimensions of the column were 3 cm in diameter and 40 cm in height.

(b) **Second column:** It was prepared to further purify fractions isolated from the first column. Stationary phase used was silica gel with mesh number of 230-400 and the mobile phase was 30% ethyl acetate/hexane. Dimensions of the column were 1.5 cm in diameter and 20 cm in height. Thin Layer Chromatography was performed on fractions isolated from first and second columns. Anisaldehyde- $\text{H}_2\text{SO}_4$  spray reagent was used for visualization.

(c) **Spectroscopic identification:** Infra-red (IR) Spectroscopy or HPLC was conducted on one of the purest fraction isolated from the second column to identify secondary metabolites.

## Research Article

### In Vitro Antioxidant Activity

To screen in vitro antioxidant activity of crude plant extracts and secondary metabolites of these ethnomedicinal plants, the following parameters were assessed spectrophotometrically:-

(i) *DPPH Radical Scavenging Activity*:-

Radical scavenging activity of leaves extracts of these plants against stable DPPH (1,1-diphenyl-2-picryl hydrazyl radical) was determined by the method of Cuendet *et al.*, (1997). (ii) *Superoxide radical scavenging activity*:-

Superoxide radical scavenging activity of plant extracts and secondary metabolites was measured by the protocol of Beauchamp and Fridovich (1971).

(iii) *Hydrogen peroxide scavenging activity*:-

This activity was assessed by following the procedure of Ruch *et al.*, (1989).

(iv) *Hydroxyl radical scavenging assay*:-

This activity was determined by following the protocol of Halliwell and Gutteridge (1981).

Antimicrobial studies were also carried out in these ethnomedicinal plants

*Test microorganisms*:

The bacterial strains studied were *Pseudomonas aeruginosa* and *Escherichia coli*. Microorganisms were maintained at 4 °C on nutrient agar slants. These test organisms were clinical isolates obtained from patients diagnosed for having bacterial infections and procured from SMS Hospital Jaipur.

### Antibacterial screening

The filter paper disc method was used for screening the extract for antibacterial activity. Standard size Whatman filter paper disc (6.0 mm diameter) was sterilized in an oven at 140°C for one hour, saturated with plant extracts such as stem and leaf and air dried at room temperature to remove any residual solvent that might interfere with the determination of activity. The discs were then placed on the surface of sterilized nutrient agar medium that had been inoculated with test bacteria (using saline solution) and air dried to remove the surface moisture. The thickness of the agar medium used was kept equal in all the petri plates. Before incubation, the petriplates were placed for one hour in cold room (5°C) to allow the diffusion of the compounds from the disc into the medium. Plates were incubated at 37°C for 20-24 hours. All the experiments were done in three replicates and the activity index was calculated.

$$\text{Activity index (A.I.)} = \frac{\text{Inhibition zone of the sample}}{\text{Inhibition zone of the standard}}$$


## RESULTS AND DISCUSSION

The ethnomedicinal plants were found to have the presence of active constituents which were responsible to cure various health ailments of the tribal people residing in that particular area

**Table 2: Antibacterial bioassay**

Pathogenic bacteria	<i>Soymida febrifuga</i>		
	ZOI of Standard	Methanolic Extract	
		Leaves ZOI	AI
<i>Escherichia coli</i>	17.6±0.28		
<i>Staphylococcus aureus</i>	19.07±0.33	30.25±0.29	1.58

Standards: Ciprofloxacin; Activity index = ZOI of test sample/ZOI of standard. Mean±SE, \*  $p < 0.05$ , \*\*  $p < 0.01$ , All columns (ZOI) are compared with standard using Tukey test after ANOVA using ezanova software



**Figure 1: Qualitative phytochemical analysis in Ethnomedicinal plants**

The results of the antimicrobial tests carried out are shown in the following tables.

**Table 3: Antibacterial bioassay**

Pathogenic bacteria	<i>Celastrus paniculata</i>									
	Standard	Methanolic Extract					Ethanolic Extract			
		Leaves		Stem			Leaves		Stem	
		ZOI	AI	ZOI	AI		ZOI	AI	ZOI	AI
<i>Escherichia coli</i>	17.6±0.28	35.33±1.45	2.00	22.66±0.66	1.28		36.66±0.66	2.08	26.66±0.88	1.51
<i>Pseudomonas aeruginosa</i>	14±0.33	11.00±0.57	0.78	10.33±0.88	0.88		14±0.57	1	12.33±0.88	0.8



**Figure 2: Zone of inhibition in *Celastrus paniculata***

# Research Article

**Table 4: Antibacterial bioassay**

Pathogenic bacteria	<i>Gmelina arborea</i>		
	Standard	Methanolic Extract	
		Leaves	
		ZOI	AI
<i>Escherichia coli</i>	17.6±0.28		
<i>Staphylococcus aureus</i>	19.07±0.33	31.00±0.29	1.62

Standards: Ciprofloxacin; Activity index = ZOI of test sample/ZOI of standard. Mean±SE, \*  $p < 0.05$ , \*\*  $p < 0.01$ , All columns (ZOI) are compared with standard using Tukey test after ANOVA using ezanova software

**Table 5: Antibacterial bioassay**

Pathogenic bacteria	<i>Gymnema sylvestris</i>				
	Standard	Methanolic Extract			
		Leaves		Stem	
		ZOI	AI	ZOI	AI
<i>Escherichia coli</i>	17.6±0.28				
<i>Staphylococcus aureus</i>	18.07±0.33	31.66±0.23	1.66	27.66±0.52	1.45

Standards: Ciprofloxacin; Activity index = ZOI of test sample/ZOI of standard. Mean±SE, \*  $p < 0.05$ , \*\*  $p < 0.01$ , All columns (ZOI) are compared with standard using Tukey test after ANOVA using ezanova software

**Table 6: Antibacterial bioassay**

Pathogenic bacteria	Peganum harmala								
	Standard	Methanolic Extract				Ethanolic Extract			
		Leaves		Stem		Leaves		Stem	
		ZOI	AI	ZOI	AI	ZOI	AI	ZOI	AI
Escherichia coli	16±0.28	7±0.57	0.41	8±1.00	0.47	5.66±0.33	0.33	-	-
Pseudomonas aeruginosa	14±0.33	6.33±0.33	0.4	-	-	-	-	-	-

Standards: Ciprofloxacin; Activity index = ZOI of test sample/ZOI of standard. Mean±SE, \*  $p < 0.05$ , \*\*  $p < 0.01$ , All columns (ZOI) are compared with standard using Tukey test after ANOVA using ezanova software

## CONCLUSION

The ethno medicinal plants used by the tribal people of south east region of Rajasthan are overexploited and hence are facing the danger of extinction therefore there is an urgent need for their conservation as they play a major role in the traditional healing system of medicine.

## REFERENCES

- Aebi H (1984). Catalase *in vitro*, In: *Method in Enzymology* by Colowick, S.P. and Kaplan, N.O. (Eds.), Academic Press, New York, **105** 121-6.  
 Ahmad I, Aqil F (2007). In vitro efficacy of bioactive extracts of 15 medicinal plants against Esbetal-producing multidrug-resistant enteric bacteria *Microbiol Research*, **162**(3) 264-275



### Research Article

- Alfermann A, Petersen M, Fuss E (2003).** Production of natural products by plant cell biotechnology: Results, problems and perspectives. In M Lamier, W Rucker, eds, *Plant Tissue Culture 100 Years Since Gottlieb Haberlandt*. Springer, New York, pp 153-166.
- Ali A, Ali M (2012).** New triterpenoids from *Morus alba* L. stem bark. *Natural Products Research*, 10.1080/14786419.2012.676547.
- Baldi A, Bisaria VS, Srivastava AK (2007).** Biotechnological approaches for the production of some promising plant-based chemotherapeutics. In: O Kayser, W Quax, eds, *Medicinal Plant Biotechnology*. Wiley, Weinheim, 117-156.
- Beauchamp C, Fridovich I (1971).** Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry*, **13**, 276-277. doi: 10.1016/0003-2697(71)90370-8.
- Cuendet M, Hostettmann K, Potterat O (1997).** Iridoid glucosides with free radical scavenging properties from *Fagraea blumei*. *Helvetica Chimica Acta*, **80**, 1144-52.
- De Luca V, St Pierre B (2000).** The cell and developmental biology of alkaloid biosynthesis. *Trends Plant Science*, 5 168-173.
- Fukai T, Hano Y, Hirakura K, Nomura T, Uzawa J (1984)** Structure of Mulberrofuran H, A Novel 2-arylbenzofuran derivative from the cultivated mulberry tree *Morus lhou* (Ser.) Koidz. *Chemical and Pharmaceutical Bulletin*, **32**(2) 808-811.
- Ghisalberti EL (2000)** *Lantana camara* L. (Verbenaceae). *Fitoterapia*, **71**(5) 467-486.
- Gulcin I, Topal F, Cakmakc R, Bilsel M, Goren AC, Erdogan U (2011)** Pomological features, nutritional quality, polyphenol content analysis and antioxidant properties of domesticated and three wild ecotype forms of raspberries (*Rubus idaeus* L.), *Journal of Food Science*, **13**(4) 585-593. doi: 10.1111/j.1750-3841.2011.02142.x.
- Halliwell B, Gutteridge JMC (1981)** Formation of thiobarbituric acid reactive substances from deoxyribose in the presence of iron salts: the role of superoxide and hydroxyl radicals. *FEBS Letters*, **13** 347-352. doi: 10.1016/0014-5793(81)80114-7. [PubMed: 6266877].
- Johnson TS, Ravishankar GA, Venkataraman LV (1990)** In vivo capsaicin production by immobilized cells and placental tissues of *Capsicum annuum* L. grown in liquid medium. *Plant Science* **70** 223-229
- Kim GN, Jang HD (2011).** Flavonol content in the water extract of the mulberry (*Morus alba* L.) leaf and their antioxidant capacities. *Journal of Food Science*, **76**(6) C869-C873.
- Kirtikar KR, Basu BD (1933).** Indian Medicinal Plants. Parabasi Press, Calcutta, India.
- Kodati DR, Burra S, Kumar Goud P (2011)** Evaluation of wound healing activity of methanolic root extract of *Plumbago zeylanica* L. in wistar albino rats. *Asian Journal of Plant Sciences and Research*, **1**(2):26-34.
- Krishnaswamy M, Purushothaman KK (1980)** Plumbagin: a study of its anticancer, antibacterial and antifungal properties. *Indian Journal of Experimental Biology*, **18**(8) 876-877.
- Malik CP, Kaur B, Verma A, Wadhvani C (2008)** Development and standardization of herbal medicines:an overview and current status.In *Herbalscures: Traditional approach*. Aavishkar Publishers, Jaipur. 473.
- Marcocci L, Maguire JJ, Droy-Lefaix MT, Packer L (1994).** The nitric oxide-scavenging properties of *Ginkgo biloba* extract EGb 761. *Biochem Biophys Res Commun* **15** 748-755.
- Marklund S, Marklund G (1974).** Involvement of superoxide anion radical in auto-oxidation of pyragallol and convenient assay for superoxide dismutase. *European Journal of Biochemistry*, **47** 469-74.
- Ming Y, Wang J, Yang J, Liu W (2011).** Chemical constituents of *Plumbago zeylanica*. *Advanced Materials Research*. **308-310** 1662-1664.
- Moron MJ, Depierr JW, Mannervik B (1979).** Levels of GSH,GR and GST activities in rat lung and liver. *Biochem Biophys Acta*, **582** 67-78.
- Obolentseva GV, Litvinenko VI, Ammosov AS (1999).** *Pharma Chemistry Journal*, **33** 24-31.
- Ohkawa H, Ohishi N, Yogi K (1979).** Assay for lipid peroxidation in chemical tissue by thiobarbituric acid reaction. *Analytical Biochemistry*, **95** 351.

**Research Article**

**Prashar R, Kumar A (1994).** Chemopreventive action of *Ocimum sanctum* on DMBA -Induced pappilomagenesis in the skin of mice. *International Journal of Pharmacy*, **33/2** 181-18.

**Preil W (2005).** A personal reflection on the use of liquid media for in vitro culture. In A Hvoslef-Eide, W Preil, eds, *Liquid Systems for in vitro Plant Propagation*. Springer, Dordecht, 1-18.

**Ravikumar VR et al. (2011).** Phytochemical and antimicrobial studies on *Plumbago zeylanica* (L) (Plumbaginaceae). *International Journal of Research In Pharmacy and Chemistry*. 1(2) 185-188.

**Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999).** Antioxidant activity applying an improved ABTS radical cation decolorisation assay. *Free Radical Biology and Medicine*, 1231-1237.

**Ruch RJ, Cheng SJ, Klaunig JE (1989).** Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea. *Carcinogen*. **13** 1003-1008. doi: 10.1093/carcin/10.6.1003.

**Sreejayan, Rao MN (1997).** Nitric oxide scavenging by curcuminoids. *Journal of Pharmacy and Pharmacology*, **49**(1)105-7.

**Vijver LMV, Lotter AP (1971).** The constituents in the roots of *Plumbago auriculata* LAM and *Plumbago zeylanica* L. responsible for antibacterial activity. *Planta Medica*, **22** 8-13.

**Wink M, Alfermann AW, Franke R, Wetterauer B, Distil M, Windhovel J, Krohn O, Fuss E, Garden H, Mohagheghzaedh A, Wildi E, Ripplinger P (2005).** Sustainable bioproduction of phytochemicals by plant in vitro cultures: anticancer agents. *Plant Genetic Resources*, **3** 90-100.